

Biogenic Amines by LCMS using Kinetex 1.7u C18 50x3.0mm

Column: Kinetex® 1.7 µm C18 100 Å, LC Column 50 x 3 mm, Ea

Dimensions: 50 x 3 mm ID

Order No: 00B-4475-Y0

Elution Type: Gradient

Eluent A: 30mM Ammonium formate

Eluent B: Acetonitrile

Gradient Profile:	Step No.	Time (min)	Pct A	Pct B
	1	0	80	20
	2	3	40	60
	3	3.1	0	100
	4	4	0	100
	5	4.1	80	20
	6	5	80	20


Flow Rate: 600

Col. Temp.: ambient

Detection: Circular Dichroism @ 0.000000000 (ambient)



Products used in this application:



Biogenic Amines by LCMS using Kinetex 1.7u C18 50x3.0mm

ANALYTES:

- 1** IQx
- 2** 8-MeIQx
- 3** IQ
- 4** 7,8-DiMeIQx
- 5** 4,8-DiMeI-Qx
- 6** MeIQ
- 7** Trp-P-2
- 8** Harman
- 9** PhIP
- 10** Norharman
- 11** AaC
- 12** MeAaC



Sample Preparation Details

for HPLC Application ID No.: 20685

Biogenic Amines by LCMS using Kinetex 1.7u C18 50x3.0mm

PRODUCT DESCRIPTION:

StrataTM-X-C 33 μ m Polymeric Strong Cation, 60 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S029-UBJ

SOLID PHASE EXTRACTION (SPE) PROCEDURE:

Note: The solvent volumes shown below are for a 60 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

Wash:

Dry:

Elute:

Final Prep and Analysis:

Inject: 0 μ L on HPLC Circular Dichroism @ 0.0000000000 (ambient)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	pKa	% Rec	%RSC (n=0)
1 IQx	0				
2 8-MeIQx	0				
3 IQ	0				
4 7,8-DiMeIQx	0				
5 4,8-DiMeI-Qx	0				
6 MeIQ	0				
7 Trp-P-2	0				
8 Harman	0				
9 PhIP	0				
10 Norharman	0				
11 AaC	0				
12 MeAaC	0				

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals.
Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com

support@phenomenex.com